Any child or teenager with symptoms that are unusual for him or her, or are persistent or unexplained, should be seen by a GP. If you are worried, make an appointment with your doctor.

Please remember any child or teenager needing urgent medical help should be taken to the nearest emergency department or dial 999.

10 children and teenagers are diagnosed with a brain tumour every week in the UK. That's more than one a day.

Early diagnosis of brain tumours can save lives.

HeadSmart is funded and promoted by The Brain Tumour Charity and run in partnership with the Children's Brain Tumour Research Centre (CBTRC) and the Royal College of Paediatrics and Child Health (RCPCH).

If you would like to talk to someone about brain tumours, please contact the Support & Info Line at The Brain Tumour Charity on:

Freephone
0808 800 0004
or email
support@thebraintumourcharity.org

headsmart.org.uk

The Brain Tumour Charity Registered Charity No. 1150054 (England and Wales) SC045081 (Scotland). CBTRC Charitable Status Inland Revenue No. X15294, RCPCH Registered Charity No. 1057744 (England and Wales) SC03829 (Scotland) 0 2017 The Brain Tumour Charity, CBTRC and RCPCH.

HEAD SMART

EARLY DIAGNOSIS OF BRAIN TUMOURS

symptoms card

This card is designed to help you know and spot the signs and symptoms of brain tumours in children and teenagers.

BABIES

UNDER 5 YEARS



Persistent/recurrent vomiting



Balance/co-ordination/walking problems



Abnormal eye movements or suspected loss of vision



Behaviour change, particularly lethargy



Fits or seizures (not with a fever)



Abnormal head position such as wry neck, head tilt or stiff neck



Increasing head circumference (crossing centiles)

If your child has one of these, see your doctor, if two or more, ask for an 'urgent referral'

CHILDREN

5 - 11 YEARS



Persistent/recurrent headache



Persistent/recurrent vomiting



Balance/co-ordination/ walking problems



Abnormal eye movements



Blurred or double vision/ loss of vision



Behaviour change



Fits or seizures



Abnormal head position such as wry neck, head tilt or stiff neck

If your child has one of these, see your doctor, if two or more, ask for an 'urgent referral'



TEENS

12 - 18 YEARS



Persistent/recurrent headache



Persistent/recurrent vomiting



Balance/co-ordination/ walking problems



Abnormal eye movements



Blurred or double vision/ loss of vision



Behaviour change



Fits or seizures



Delayed or arrested puberty



